arXiv:1304.2856v1 [cond-mat.str-el] 10 Apr 2013

Single-mode approximation for rotational symmetry broken quantum Hall states

Rui-Zhi Qiu[1,2], Zi-Xiang Hu[3], and Xin Wan[1]
[1]*Zhejiang Institute of Modern Physics, Zhejiang University, Hangzhou 310027, P.R. China*
[2]*Science and Technology on Surface Physics and Chemistry Laboratory,*
P.O. Box 718-35, Mianyang 621907, Sichuan, P.R. China and
[3]*Department of Physics, Chongqing University, Chongqing 400044, P.R.China*

(Dated: April 11, 2013)

A topological phase can often be represented by a corresponding wavefunction (exact eigenstate of a model Hamiltonian) that has a higher underlying symmetry than necessary. When the symmetry is explicitly broken in the Hamiltonian, the model wavefunction fails to account for the change due to the lack of a variational parameter. Here we exemplify the case by an integer quantum Hall system with anisotropic interaction. We demonstrate the recovery of the variational parameter in a single-mode approximation, which is consistent with the recently proposed geometric consideration of the quantum Hall phases.

Introduction. The integer quantum Hall (IQH) effect is now regarded as a simple class of topological insulators with broken time-reversal symmetry. A system in such a phase has invariant properties of its ground state wavefunction under smooth deformations of the Hamiltonian. This, of course, does not mean that the ground state wavefunction itself is invariant. For convenience, we often choose to study the system with continuous symmetries in translation and/or rotation in which the ground state wavefunction is relatively simple, although these symmetries are unnecessary for the study of the topological properties. In fact, such a choice can sometimes hide important physics.

Consider an IQH system in the disk geometry with rotational symmetry. Electrons in a polarized lowest Landau level (LLL) fill up ring-shaped orbitals $\phi_m(z) = z^m e^{-|z|^2/4}$, where $z = x + iy$ is the complex coordinate on the plane. The trivial many-body ground state is the Slater determinant of the orbitals with the lowest possible m

$$\Phi_0(z_1, ..., z_N) = \left[\prod_{i<j}(z_i - z_j)\right] e^{-\sum_i |z_i|^2/4}, \qquad (1)$$

which is the well known Vandermonde determinant multiplied by the ubiquitous Gaussian factor. The wavefunction, which vanishes when two electrons coincide with each other, manifests the Pauli exclusion principle. The trivial many-body state is known to be the ground state even in the presence of interaction.

This prompts an interesting question: Can effects of interaction be explicitly manifested in the robust IQH state? If yes, how do they modify the ground state wavefunction variationally? In fact, to construct a variational wavefunction for correlated systems from its noninteracting counterpart is a long-standing problem [1]. Suppose we decompose a many-body Hamiltonian to be $H = H_0 + H_1$. We assume that we know the ground state $|\Phi_0\rangle$ of H_0, which describes, for the present discussion, the noninteracting part of the total Hamiltonian. H_1 describes the interparticle interaction. Within the single-mode approximation, which replaces the effect of H_1 by a single, mean excitation energy ω_0, we can write down an ansatz wavefunction $|\Psi_0\rangle = e^{-H_1/\omega_0}|\Phi_0\rangle$ for the ground state of H. Here, ω_0 can be tuned as a variational parameter to minimize the ground state energy. A well-known example is the Gutzwiller's wavefunction [2], which accounts for the electron correlation in the on-site Hubbard model. An improved independent-mode approximation can lead to Jastrow's ansatz for a trial ground-state wavefunction

$$\Psi_0(\mathbf{r}_1, ..., \mathbf{r}_N) = e^{\sum_{ij} f(\mathbf{r}_i - \mathbf{r}_j)} \Phi_0(\mathbf{r}_1, ..., \mathbf{r}_N), \qquad (2)$$

where the Jastrow function $f(\mathbf{r} - \mathbf{r}')$ is determined by energy minimization [1]. According to this description, the Jastrow function for the IQH state simply vanishes despite the interparticle interaction.

This, of course, is counterintuitive. In fact, the absence of the Jastrow factor for the isotropic IQH state, which can be regarded as a special case of the Laughlin state at filling fraction $\nu = 1$, has already been questioned in the context of the geometric description of fractional quantum Hall (FQH) states [4, 5]. Haldane [4] argued that guiding center metric should be regarded as a variational parameter in the generalized family of Laughlin trial wavefunctions. Explicit construction of the family of wavefunctions [5] confirms that the isotropic Laughlin wavefunction is the variational ground state for an isotropic interparticle interaction. In contrast, the dipole-dipole interaction, as occurs in rotating dipolar cold atomic systems, leads to an anisotropic variational ground state [5]. On a torus geometry without boundary, one can calibrate the wavefunction anisotropy or the guiding center metric by exploring static structure factor and pair correlation function [6, 7].

The geometric description of quantum Hall states echoes the single-mode (independent-mode) approximation in that both quest for a (set of) variational parameter(s) that accounts for the correlation of electrons. It is, therefore, conceivable that for the IQH effect the two descriptions may be unified. It should be pointed out that the variational parameter also emerged from the discussion of the stress response to an applied field by Read and Rezayi [8] for two-dimensional electrons in a uniform magnetic field. Although the authors there turned to reveal the illuminating relation of the Hall viscosity to the mean orbital spin per particle, their metric-deformed single-particle basis is sufficient to set up the stage for a vari-



FIG. 1: (Color online) The mean total angular momentum of the ground state $\overline{M} = \langle \Psi^{(N)}(\alpha,\theta) | L^z | \Psi^{(N)}(\alpha,\theta) \rangle$ (in units of $\hbar$) in a system with $N = 6$ particles for various confining potential strength α and tilt angle θ. The blue region to the top and right of the orange dashed line represents the anisotropic IQH state, while the rest is loosely referred to as the FQH regime, with the brown region in the lower left corner representing the $\nu = 1/3$ Laughlin state. The IQH regime is separated into the perturbative and nonperturbative IQH regimes, depending on whether Eq. (9) is a good description of the ground state or not. Numerical analysis suggests that $\overline{M} - M_0 = 0.5$ can be used to define the boundary between the two IQH regimes.

ational IQH wavefunction [5, 8]

$$\Phi_0^{\gamma}(z_1, ..., z_N) = e^{-\gamma \sum_i z_i^2/4} \Phi_0(z_1, ..., z_N), \tag{3}$$

where γ parametrizes the metric change or anisotropy. This apparently differs from Eq. (2), but we will argue they are consistent, at least for the IQH state with dipole-dipole interaction.

In this Letter we take a pedagogical approach to construct the variational ground-state wavefunction from numerical diagonalization of a microscopic system with dipole-dipole interaction. We find that in the perturbative regime the realistic ground state can be written as the isotropic IQH ground state multiplied by the sum of the leading terms in a geometric series, which can be cast in an exponential form much like the Jastrow factor in Eq. (2). The Jastrow factor can be understood as arising, in the single-mode approximation, from an anisotropic quadrupolar interaction. Up to a center-of-mass contribution not generated by the dipole-dipole interaction, we show that the wavefunction agrees with the Bogoliubov transformation of the guiding center coordinates in the isotropic wavefunction, or Eq. (3).

Model. As proposed in Ref. 9, the fast rotating quasi-two-dimensional gas of polarized fermionic dipoles serves as an ideal arena for the present study. The Larmor theorem states that the fast rotation is equivalent to a high magnetic field for the fermions. Since the p-wave interaction for the polarized fermions is typically small unless in the resonance regime, the only significant interaction in the system is the dipole-dipole interaction. Therefore, we set up a quantum Hall system with anisotropic dipole-dipole interaction as described by the following Hamiltonian



FIG. 2: (Color online) Low-energy excitations for the isotropic IQH state with $\alpha = 0.1$ and $\theta = 0°$. The blue thick levels can be identified as Φ_0 [Eq. (1)], Φ_2^2 [Eq. (7)], and Φ_4^2 [Eq. (8)]. Note that their excitation energies ΔE scale with their momenta ΔM. These states are the main components of the anisotropic IQH ground state for $\theta \neq 0°$.

$$H = \alpha L^z + \frac{1}{2} \sum_{m1,m2,m3,m4} V_{1234}(\theta) f_{m1}^\dagger f_{m2}^\dagger f_{m4} f_{m3}, \tag{4}$$

where $f_m^\dagger$ creates a fermion in the state ϕ_m and $L^z = \sum_m m f_m^\dagger f_m$ is the z-component of the total angular momentum. The details of the anisotropic dipole-dipole interaction matrix elements $V_{1234}(\theta)$ are explained in Ref. [9]. The isotropic confining potential strength α can be tuned by the rotation frequency and the tilt angle θ by applied electric and magnetic field. We explore, by exact diagonalization, the N-particle ground state and compute its mean total angular momentum (in units of $\hbar$)

$$\overline{M} \equiv \left\langle \Psi^{(N)}(\alpha,\theta) \middle| L^z \middle| \Psi^{(N)}(\alpha,\theta) \right\rangle, \tag{5}$$

where $\Psi^{(N)}(\alpha,\theta)$ denotes the ground state wave function for given parameters α and θ.

As illustrated in Fig. 1, with isotropic dipole-dipole interaction in the plane of motion, i.e. for $\theta = 0°$, the total angular moment is a good quantum number and can be matched to that of the IQH state $M_0 = N(N-1)/2$ for $\alpha > 0.085$ and $M_L = 3N(N-1)/2$ the $\nu = 1/3$ Laughlin state for $\alpha < 0.018$. Note that we only show $\alpha > 0.01$ to avoid the unwanted complication due to weak confinement and finite system size. As θ increases, the rotational symmetry is explicitly broken and $\overline{M}$ increases from M_0 to reflect the anisotropy of the interaction. We identify the extended blue region to the up and right of the curved orange dashed line in Fig. 1 to be the *anisotropic IQH phase* with $\overline{M} = M_0 + O(1)$, which has the maximum density in the interior of the oval-shaped



FIG. 3: (Color online) Overlap analysis of $1 - |\langle\Psi^{(6)}(\alpha,\theta)|\Phi_0\rangle|^2$, $|\langle\Psi^{(6)}(\alpha,\theta)|\Phi_2^2\rangle|^2$, and $|\langle\Psi^{(6)}(\alpha,\theta)|\Phi_4^2\rangle|^2$ for (a) various θ at a fixed $\alpha = 0.1$ and (b) various α at a fixed $\theta = 70^\circ$. The overlap analysis suggests that $\Psi^{(6)}(\alpha = 0.1, \theta)$ is mostly composed of Φ_0, Φ_2^2, and Φ_4^2. In (b) we also plot $|\langle\Psi^{(6)}(\alpha,\theta)|\Phi_4^2\rangle|^2$ calculated from Eq. (10), which only deviates from the direct overlap calculation for weak confinement $\alpha < 0.03$.

droplet [9]. As labeled in Fig. 1, the IQH phase can be further separated into perturbative and nonperturbative regimes to be explained later. The rest of the parameter space is loosely referred to as the *FQH regime* [9].

Ground-state wavefunction. The isotropic IQH ground state (in a harmonic trap) is the maximum density droplet of electrons in the LLL. One expects that in the presence of small anisotropic interaction the IQH state can only be perturbed at the edge. Hence the anisotropic IQH state is expected to be the superposition of the isotropic IQH ground state and its edge states, whose spectrum is shown in Fig. 2 for $\alpha = 0.1$. Due to the nature of the dipole-dipole interaction, only edge states with an angular momentum difference from the ground state by an integral multiple of 2 can be generated. Therefore, the leading correction must come from either one of the following states or both,

$$\Phi_2^1 = \left[\sum_i z_i\right]^2 \prod_{i<j}(z_i - z_j)e^{-\sum_i |z_i|^2/4}, \qquad (6)$$

$$\Phi_2^2 = \left[\sum_{i<j}(z_i - z_j)^2\right] \prod_{i<j}(z_i - z_j)e^{-\sum_i |z_i|^2/4}, \qquad (7)$$

which are unnormalized but orthogonal to each other in the thermodynamic limit. The nomralization factor can be worked out using the algebra of symmetric polynomials. Numerical analysis reveals that the ground state has vanishingly small overlap with Φ_2^1, but substantially large overlap with Φ_2^2. In fact, both Φ_2^2 and Φ_0 are known to be the eigenstates of the projected isotropic interaction in the lowest Landau level [10], which are indicated in the low-lying energy levels for the corresponding isotropic interaction in Fig. 2. In the figure we also highlight a third energy level, which corresponds to

θ	0°	20°	40°	60°	80°
$\lvert\langle\Psi^{(6)}(0.1,\theta)\lvert\Psi_0\rangle\rvert$	1.0000	0.9994	0.9945	0.9865	0.9809
$\lvert\langle\Psi^{(6)}(0.1,\theta)\lvert\Psi_2^2\rangle\rvert$	0.0000	0.0342	0.1042	0.1628	0.1927
$\lvert\langle\Psi^{(6)}(0.1,\theta)\lvert\Psi_4^2\rangle\rvert$	0.0000	0.0008	0.0074	0.0183	0.0258

TABLE I: Overlaps of the ground state $\Psi^{(N)}(\alpha = 0.1, \theta)$ obtained by exact diagonalization of an $N = 6$ system and the wavefunctions Φ_0, Φ_2^2, and Φ_4^2.

$$\Phi_4^2 = \left[\sum_{i<j}(z_i - z_j)^2\right]^2 \prod_{i<j}(z_i - z_j)e^{-\sum_i |z_i|^2/4}. \qquad (8)$$

We should emphasize that the labeled edge excitations in Fig. 2 for realistic interaction are not exactly the model wavefunctions, i.e., specified by Eqs. (7) and (8), respectively, but the corresponding overlaps are close to unity. For example, in an $N = 6$ system with isotropic dipole-dipole interaction, the overlap of the corresponding state with Φ_4^2 is 99.85%. From the form of the wavefunctions, as well as the linear energy-momentum relation, of the states Φ_2^2 and Φ_4^2, the latter is clearly the corresponding two-boson excitation of the former.

First, we focus on the regime with $\overline{M} - M_0 \ll 1$, in which these edge-state wavefunctions are sufficient for the analysis of anisotropic IQH states. We choose $\alpha = 0.1$ (the top boundary in Fig. 1) and calculate the overlaps of the ground state $\Psi^{(N)}(\alpha,\theta)$ with $N = 6$ obtained by exact diagonalization and the wavefunctions Φ_0, Φ_2^2, and Φ_4^2. In Figure 3(a) we plot $1 - |\langle\Psi^{(6)}(\alpha,\theta)|\Phi_0\rangle|^2$, $|\langle\Psi^{(6)}(\alpha,\theta)|\Phi_2^2\rangle|^2$, and $|\langle\Psi^{(6)}(\alpha,\theta)|\Phi_4^2\rangle|^2$ as a function of θ. For convenience, we also list the values of overlap at selected angles θ in Table I. As θ increases, the ground state evolves from Φ_0 to a weighted sum of Φ_0 and Φ_2^2, and eventually to a weighted sum of Φ_0, Φ_2^2, and Φ_4^2. At $\alpha = 0.1$ the three states together exhaust $\Psi^{(N)}(\alpha,\theta)$, e.g. up to 99.996% at $\theta = 80^\circ$.

Not surprisingly, the overlap with the higher-momentum Φ_4^2 is much smaller than that with Φ_2^2. A question arises whether the former can be related to the latter in the perturbative regime. Given the general knowledge of trial wavefunctions for correlated systems, we attemped to fit the numerical wavefunction by

$$\Psi_\gamma = e^{-\gamma\sum_{i<j}(z_i - z_j)^2}\Phi_0, \qquad (9)$$

of which the exponential factor can be expanded to include Φ_0, Φ_2^2, and Φ_4^2 as the leading terms. If this is a good trial wavefunction, we expect

$$\left|\langle\Psi^{(N)}|\Phi_4^2\rangle\right| = \sqrt{\frac{N^2+1}{2(N^2-1)}}\left|\langle\Psi^{(N)}|\Phi_2^2\rangle\right|^2. \qquad (10)$$

For $N = 6$ the right side can be evaluated to be 0.0009, 0.0079, 0.0193, and 0.0270 for $\theta = 20^\circ$, 40°, 60°, and

80°, respectively. The calculated values are in good agreement with the tabulated ones in Table I. This regime, with Eq. (9) as a good variational wavefunction, is dubbed as the *perturbative IQH regime*. We point out that, in the perturbative regime, the overlap with the three-boson excitation $\Phi_6^2 = [\sum_{i<j}(z_i - z_j)^2]^3\Phi_0$ is vanishingly small.

Figure 3(b) repeats the overlap analysis for various α at a fixed $\theta = 70°$. For comparison, we also plot $|\langle\Psi^{(6)}(\alpha,\theta)|\Phi_4^2\rangle|^2$ calculated from Eq. (10). We find that $\Psi^{(N)}(\alpha,\theta)$ is still, to a good approximation, exhausted by Φ_0, Φ_2^2, and Φ_4^2. However, $|\langle\Psi^{(6)}(\alpha,\theta)|\Phi_4^2\rangle|^2$ calculated from Eq. (10) and by the overlap analysis deviate from each other for $\alpha < 0.03$. The onset of the deviation is almost independent of θ in the anisotropic IQH phase and coincides well with $\overline{M} - M_0 = 0.5$. Therefore, we dubbed the regime with larger $\overline{M}$ the *nonperturbative IQH regime*, as illustrated in Fig. 1, in which Eq. (9) is insufficient in describing the true ground state precisely.

The form of Eq. (9) exemplifies the independent-mode approximation in Eq. (2) in the IQH context. In particular, since Φ_4^2 is the two-boson excitation of Φ_2^2, the wavefunction Eq. (9) is a single-mode approximation for the anisotropic IQH state. As discussed in the introduction to the single-mode approximation, the wavefunction Eq. (9) suggests one can extract, under the single-mode approximation, a quadrupolar interaction $V(z_i, z_j) = (z_i - z_j)^2$ from the anisotropic interaction. More formal formulations require us to write down a real interaction $V(z_i, z_j) + V(\bar{z}_i, \bar{z}_j)$, whose antiholomorphic contribution drops out in the LLL projection. Not surprisingly, the real form faithfully reproduces the angular dependence of the anisotropic component of the dipole-dipole interaction.

Discussions. The consideration of the broken rotational invariance in the quantum Hall context appeared first in the alternative discussion on the effect of anisotropic effective mass tensor [11]. Very recently the joint effects of both anisotropic mass and interaction surfaced in the geometric description of quantum Hall states [4], of which the emergence of the Jastrow factor can be understood as a special example. Haldane [4] pointed out that the conventional understanding of the Laughlin wavefunctions is not complete; in his novel geometric description the original Laughlin wavefunction is simply a member of a family of Laughlin states, parameterized by a hidden (continuous) geometric degree of freedom. The family of the Laughlin states, with the geometric factor as a variational parameter, potentially provides a better description of the FQH effect in the presence of either anisotropic effective mass or anisotropic interaction, which are present in real materials. The present work also confirms that even in the IQH context the geometric description is necessary and we uncovered the variational parameter γ by our numerical anatomy of the ground state wavefunction.

The family of the variational wavefunction can be constructed alternatively by squeezing the LL orbitals. In the symmetric gauge one can introduce a unimodular (or Bogliubov) transformation

$$\begin{pmatrix} z \\ \bar{z} \end{pmatrix} \rightarrow \frac{1}{\sqrt{1-|\gamma|^2}} \begin{pmatrix} 1 & \gamma^* \\ \gamma & 1 \end{pmatrix} \begin{pmatrix} z \\ \bar{z} \end{pmatrix}. \tag{11}$$

of the LL wavefunctions to generate a set of anisotropic single-particle basis states [9]. As a result, we find that the $\nu = 1$ IQH state (in the absence of LL mixing) becomes Eq. (3). Note that the extra term in the exponential, which arises from the modification of single-particle basis, will be ubiquitous for all FQH wavefunctions. The unimodular transformation approach also appeared in the consideration of Read and Rezayi [8] on the geometric aspects of the uniform deformation of quantum Hall states in the context of Hall viscocity, which is the nondissipative transport coefficient that describes the stress in the adiabatic response to a strain. The authors [8] pointed out that the extra term in the exponential corresponds to a quadrupolar harmonic perturbation in the plasma mapping for the Laughlin state [12].

Apparently, Eq. (3) differs from Eq. (9). But the two forms can be related by noticing

$$\sum_i z_i^2 = \frac{1}{N}\left[\sum_i z_i\right]^2 + \frac{1}{N}\left[\sum_{i<j}(z_i - z_j)^2\right]. \tag{12}$$

The difference $e^{-\gamma(\sum_i z_i)^2/N}$, when expanded for small γ, involves Φ_2^1 and its descendents, which are not generated by the dipole-dipole interaction, as we discussed in the wavefunction analysis. Therefore, the unimodularly transformed wavefunction for the IQH case is consistent with the single-mode approximation of the anisotropic quadrupolar interaction.

The explicit construction of the family of variational wavefunctions in the FQH case was discussed by two of the authors and collaborators. For concreteness, the anisotropic Langhlin state at $\nu = 1/3$ is

$$\prod_{i<j}\left[z_i - z_j + 2\gamma^*(\tilde{\partial}_{z_i} - \tilde{\partial}_{z_j})\right]^3 e^{-\sum_i \gamma z_i^2/4}, \tag{13}$$

where we omit the usual Gaussian factor for simplicity; we use $\tilde{\partial}$ to emphasize that the partial derivatives are not operated on the (omitted) Gaussian factor. The wavefunction is simply the unimodular transformation of the isotropic Laughlin wavefunction projected to the LLL. For small γ the exponential, shared by the IQH wavefunction, generates edge excitations, i.e. density deformation at the edge. The Jastrow product describes the deformation of the correlation between particles, i.e. the shape of the correlation hole. As a result of the unimodular transformation construction, the two kinds of deformations are controlled by the same (complex) anisotropy parameter. For a realistic edge, the confining potential is expected to squeeze the shape of the correlation hole locally near the edge, which may offers an alternative understanding of FQH edge physics in the geometric description [13].

In summary, we find that the anisotropic IQH phase due to interparticle dipole-dipole interaction can be separated into the perturbative and nonperturbative regimes, based on the total angular momentum deviation from its isotropic value. In the perturbative regime, we show that a variational wavefunction can be constructed in the single-mode approximation as in the generic treatment for correlated systems. We further argue that the single-mode approximation for the anisotropic interaction is consistent with the geometric description for the anisotropic IQH phase.

XW acknowledges the illuminating discussions with Peter Fulde and Duncan Haldane. This work was supported by the 973 Program under Project Nos. 2009CB929101 and 2012CB927404, the NSFC Project Nos. 11174246 and 11274403, and the Discipline Development Fund Project No. ZDXKFZ201206.

[1] P. Fulde, *Electron Correlations in Molecules and Solids*, 3rd ed. (Springer, Berlin, Heidelberg 1995) p. 100.
[2] M. C. Gutzwiller, Phys. Rev. Lett. **10**, 159 (1963).
[3] R. Jastrow, Phys. Rev. **98**, 1479 (1955).
[4] F. D. M. Haldane, Phys. Rev. Lett. 107, 116801 (2011).
[5] R.-Z. Qiu, F. D. M. Haldane, X. Wan, K. Yang, and S. Yi, Phys. Rev. B **85**, 115308 (2012).
[6] B. Yang, Z. Papić, E. H. Rezayi, R. N. Bhatt, and F. D. M. Haldane, Phys. Rev. B 85, 165318 (2012).
[7] H. Wang, R. Narayanan, X. Wan, and F.-C. Zhang, Phys. Rev. B 86, 035122 (2012).
[8] N. Read and E. H. Rezayi, Phys. Rev. B 84, 085316 (2011).
[9] R.-Z. Qiu, S.-P. Kou, Z.-X. Hu, X. Wan, and S. Yi, Phys. Rev. A **83**, 063633 (2011).
[10] S. A. Trugman and S. Kivelson, Phys. Rev. B **31**, 5280 (1985).
[11] F. D. M. Haldane, in *The Quantum Hall Effect*, eds. R. E. Prange and S. M. Girvin, 2nd ed. (Springer, New York 1990) p. 309.
[12] P. Di Francesco, M. Gaudin, C. Itzykson, and F. Lesage, Int. J. Mod. Phys. A9, 4257 (1994).
[13] F. D. M. Haldane, private communication.